arXiv:1411.3631v1 [math.DG] 13 Nov 2014

CURVATURE ESTIMATES FOR FOUR-DIMENSIONAL GRADIENT STEADY RICCI SOLITONS

HUAI-DONG CAO AND XIN CUI

Abstract. In this paper, we derive certain curvature estimates for 4-dimensional gradient steady Ricci solitons either with positive Ricci curvature or with scalar curvature decay $\lim_{x\to\infty} R(x) = 0$.

1. The results

A complete Riemannian metric g_{ij} on a smooth manifold M^n is called a *gradient steady Ricci soliton* if there exists a smooth function f on M^n such that the Ricci tensor R_{ij} of the metric g_{ij} satisfies the equation

$$R_{ij} + \nabla_i \nabla_j f = 0. \tag{1.1}$$

The function f is called a *potential function* of the gradient steady soliton. Clearly, when f is a constant the gradient steady Ricci soliton (M^n, g_{ij}, f) is simply a Ricci-flat manifold. Gradient steady solitons play an important role in Hamilton's Ricci flow, as they correspond to translating solutions, and often arise as Type II singularity models. Thus one is interested in possibly classifying them or understanding their geometry.

It turns out that compact steady solitons must be Ricci-flat. In dimension $n = 2$, Hamilton [21] discovered the first example of a complete noncompact gradient steady soliton on $\mathbb{R}^2$, called the *cigar soliton*, where the metric is given by

$$ds^2 = \frac{dx^2 + dy^2}{1 + x^2 + y^2}.$$

The cigar soliton has potential function $f = -\log(1 + x^2 + y^2)$, positive curvature $R = 4e^f$, and is asymptotic to cylinder at infinity. Furthermore, Hamilton [21] showed that the only complete steady soliton on a two-dimensional manifold with bounded (scalar) curvature R which assumes its maximum at an origin is, up to scaling, the cigar soliton. For $n \geq 3$, Bryant [3] proved that there exists, up to scaling, a unique complete rotationally symmetric gradient Ricci soliton on $\mathbb{R}^n$, see, e.g., Chow et al. [14] for details. The Bryant soliton has positive sectional curvature, linear curvature decay and volume growth of geodesic balls $B(0, r)$ on the order of $r^{(n+1)/2}$. In the Kähler case, Cao [5] constructed a complete $U(m)$-invariant gradient steady Kähler-Ricci soliton on $\mathbb{C}^m$, for $m \geq 2$, with positive sectional curvature. It has volume growth on the order of r^m and also linear curvature decay. Note that in each of these three examples, maximum of the scalar curvature is attained at the origin. One can find additional examples of steady solitons, e.g., in [23, 19, 16, 17, 4] etc; see also [6] and the references therein.

In dimension $n = 3$, Perelman [27] claimed that the Bryant soliton is the only complete noncompact, κ-noncollapsed, gradient steady soliton with positive sectional curvature. Recently, Brendle has affirmed this conjecture of Perelman (see [1]; and also [2] for an extension to the higher dimensional case). On the other hand, for $n \geq 4$, Cao-Chen [8] and Catino-Mantegazza [11] proved independently, and using different methods, that any n-dimensional complete noncompact locally conformally flat gradient steady Ricci soliton (M^n, g_{ij}, f) is either flat or isometric to the Bryant soliton (the method of Cao-Chen [8] also applies to the case of dimension $n = 3$). In addition, Bach-flat gradient steady solitons (with positive Ricci curvature) for all $n \geq 3$ [7] and half-conformally flat ones for $n = 4$ [13] have been classified respectively.

Inspired by the very recent work of Munteanu-Wang [25], in this paper we study curvature estimates of 4-dimensional complete noncompact gradient steady solitons. In [25], Munteanu and Wang made an important observation that the curvature tensor of a four-dimensional gradient Ricci soliton (M^4, g_{ij}, f) can be estimated in terms of the potential function f, the Ricci tensor and its first derivates. In addition, the (optimal) asymptotic quadratic growth property of the potential function f proved in [9], as well as a key scalar curvature lower bound $R \geq c/f$ shown in [15] are crucial in their work. While gradient steady Ricci solitons in general don't share these two special features (cf. [31, 24, 30] and [15, 20]), some of the arguments in [25] can still be adapted to prove certain curvature estimates for two classes of gradient steady solitons. Our main results are:

Theorem 1.1. *Let (M^4, g_{ij}, f) be a complete noncompact 4-dimensional gradient steady Ricci soliton with positive Ricci curvature $Ric > 0$ such that the scalar curvature R attains its maximum at some point $x_0 \in M^4$. Then, (M^4, g_{ij}) has bounded Riemann curvature tensor, i.e.*

$$\sup_{x \in M} |Rm| \leq C$$

for some constant $C > 0$. Suppose in addition R has at most linear decay, then

$$\sup_{x \in M} \frac{|Rm|}{R} \leq C.$$

Theorem 1.2. *Let (M^4, g_{ij}, f), which is not Ricci-flat, be a complete noncompact 4-dimensional gradient steady Ricci soliton. If $\lim_{x \to \infty} R(x) = 0$, then, for each $0 < a < 1$, there exists a constant $C > 0$ such that*

$$|Ric|^2 \leq CR^a \qquad \text{and} \qquad \sup_{x \in M} |Rm| \leq C.$$

Suppose in addition R has at most polynomial decay. Then, for each $0 < a < 1$, there exists a constant $C > 0$ such that

$$|Rm|^2 \leq CR^a.$$

Remark 1.1. We have learned that the first part of Theorem 1.2 was also known to O. Munteanu and J. Wang.

Acknowledgments. We are grateful to Ovidiu Munteanu and Jiaping Wang for sending us their paper [25], and its early version in July 2014, which motivated us to consider curvature estimates for 4D steady solitons. The first author also would like to thank Ovidiu Munteanu for very helpful discussions; part of the work

was carried out when he was visiting University of Macau, partially support by FDCT/016/2013/A1 and RDG010, during summer 2014.

2. Preliminaries

In this section, we recall some basic facts and collect several known results about gradient steady solitons. Throughout the rest of the paper, we denote by

$$Rm = \{R_{ijkl}\}, \quad Ric = \{R_{ik}\}, \quad R$$

the Riemann curvature tensor, the Ricci tensor, and the scalar curvature of the metric g_{ij} respectively.

Lemma 2.1. (Hamilton [22]**)** *Let (M^n, g_{ij}, f) be a complete gradient stead soliton satisfying Eq. (1.1). Then*

$$R = -\Delta f, \tag{2.1}$$

$$\nabla_i R = 2R_{ij}\nabla_j f, \tag{2.2}$$

$$R + |\nabla f|^2 = C_0 \tag{2.3}$$

for some constant C_0.

Next we need the following useful result by B.-L. Chen [12].

Proposition 2.1. *Let $g_{ij}(t)$ be a complete ancient solution to the Ricci flow on a noncompact manifold M^n. Then the scalar curvature R of $g_{ij}(t)$ is nonnegative for all t.*

As an immediate corollary, we have

Lemma 2.2. *Let (M^n, g_{ij}, f) be a complete gradient steady soliton. Then it has nonnegative scalar curvature $R \geq 0$.*

Remark 2.1. In fact, by Proposition 3.2 in [28], either $R > 0$ or (M^n, g_{ij}) is Ricci flat.

It follows from Lemma 2.2 that the constant C_0 in (2.3) is positive whenever f is a non-constant function (i.e., the steady soliton is non-trivial). By a suitable scaling of the metric g_{ij}, we can normalize $C_0 = 1$ so that

$$R + |\nabla f|^2 = 1. \tag{2.4}$$

In the rest of the paper, we shall always assume this normalization (2.4).

Combining (2.1) and (2.4), we obtain $-\Delta f + |\nabla f|^2 = 1$. Thus, setting $F = -f$, we have

$$\Delta_f F = 1, \tag{2.5}$$

where

$$\Delta_f =: \Delta - \nabla f \cdot \nabla. \tag{2.6}$$

For gradient steady solitons with positive Ricci curvature $Ric > 0$, we have

Proposition 2.2. *Let (M^n, g_{ij}, f) be a complete noncompact gradient steady soliton with positive Ricci curvature $Ric > 0$ such that the scalar curvature R attains its maximum $R_{max} = 1$ at some origin $x_0 \in M^n$. Then, there exist some constants $0 < c_1 \leq 1$ and $c_2 > 0$ such that $F = -f$ satisfies the estimates*

$$c_1 r(x) - c_2 \leq F(x) \leq r(x) + |F(x_0)|, \tag{2.7}$$

where $r(x) = d(x_0, x)$ is the distance function from x_0.

Remark 2.2. In (2.7), only the lower bound on F requires the assumptions on Ric and R.

Note that, under the assumption in Proposition 2.2, $F(x)$ is proportional to the distance function $r(x) = d(x_0, x)$ from above and below. Throughout the paper, we denote

$$\begin{aligned} D(t) &= \{x \in M : F(x) \le t\}, \\ B(t) = B(x_0, t) &= \{x \in M : d(x_0, x) \le t\}. \end{aligned}$$

We also collect several differential equalities on R, Ric and Rm which are special cases of evolution equations of the curvature tensors under the Ricci flow:

Lemma 2.3. *Let (M^n, g_{ij}, f) be a complete gradient steady soliton satisfying Eq. (1.1). Then, we have*

$$\begin{aligned} \Delta_f R &= -2|Ric|^2, \\ \Delta_f Ric &= -2R_{ijkl}R_{jl}, \\ \Delta_f Rm &= Rm * Rm, \end{aligned}$$

where the RHS of the last equation denotes (a finite number of) terms involving quadratics in Rm.

Based on Lemma 2.3, one can easily derive the following inequalities similar to [25]:

Lemma 2.4. *Let (M^n, g_{ij}, f) be a complete gradient steady soliton satisfying Eq. (1.1). Then*

$$\begin{aligned} \Delta_f |Ric|^2 &\ge 2|\nabla Ric|^2 - 4|Rm||Ric|^2, \\ \Delta_f |Rm| &\ge -c|Rm|^2, \\ \Delta_f |Rm|^2 &\ge 2|\nabla Rm|^2 - C|Rm|^3. \end{aligned}$$

Here $c > 0$ is some universal constant depending only on the dimension n.

Remark 2.3. To derive the second differential inequality, one needs to use the Kato inequality $|\nabla |Rm|| \le |\nabla Rm|$ as shown in [25].

3. 4D gradient steady solitons with positive Ricci curvature

First of all, we need the following key fact, valid for 4-dimensional gradient steady Ricci solitons in general, due to Munteanu and Wang [25].

Lemma 3.1. (Munteanu-Wang [25]**)** *Let (M^4, g_{ij}, f) be a complete noncompact gradient steady soliton satisfying (1.1). Then, there exists some universal constant $c > 0$ such that*

$$|Rm| \le c\left(\frac{|\nabla Ric|}{|\nabla f|} + \frac{|Ric|^2}{|\nabla f|^2} + |Ric|\right).$$

Proof. This follows from the same arguments as in the proof of Proposition 1.1 of [25], but using Lemma 2.4 instead and without replacing $|\nabla f|^2$ by f in their argument.

□

Proposition 3.1. *Let (M^4, g_{ij}, f) be a complete noncompact gradient steady soliton with positive Ricci curvature and R attains maximum. Then, there exists some constant $C > 0$, depending on the constant c_1 in (2.7), such that outside a compact set,*

$$|Rm| \leq C(|\nabla Ric| + |Ric|^2 + |Ric|).$$

Proof. This easily follows from Lemma 3.1 and the following fact shown by Cao-Chen [8],

$$|\nabla f|^2 \geq c_1 > 0. \tag{3.1}$$

□

Remark 3.1. Note that, combining (3.1) with (2.4) and Lemma 2.2, we have

$$0 < c_1 \leq |\nabla F|^2 = |\nabla f|^2 \leq 1. \tag{3.2}$$

Now we are ready to prove the first main result of our paper.

Theorem 3.1. *Let (M^4, g_{ij}, f) be a complete noncompact gradient steady soliton with positive Ricci curvature $Ric > 0$ such that R attains its maximum at some point $x_0 \in M^4$. Then, there exists some constant $C > 0$, depending on c_1 in (2.7), such that*

$$\sup_{x \in M} |Rm| \leq C.$$

Proof. First of all, from (2.4), we have $R \leq 1$. Hence, since $Ric > 0$, it follows that

$$0 < |Ric| \leq R \leq 1. \tag{3.3}$$

Thus, by Proposition 3.1 and (3.3), we see that

$$|\nabla Ric|^2 \geq \frac{1}{2C^2}|Rm|^2 - (|Ric|^2 + |Ric|)^2 \geq \frac{1}{2C^2}|Rm|^2 - 4. \tag{3.4}$$

Using the first two inequalities in Lemma 2.4, we obtain

$$\Delta_f(|Rm| + \lambda|Ric|^2) \geq -C|Rm|^2 + 2\lambda(|\nabla Ric|^2 - 2|Rm||Ric|^2). \tag{3.5}$$

By (3.3), (3.4), and picking constant $\lambda > 0$ sufficiently large (depending on the constant C in Proposition 3.1, hence on c_1), it follows that

$$\Delta_f(|Rm| + \lambda|Ric|^2) \geq 2|Rm|^2 - 4\lambda|Rm| - C' \geq (|Rm| + \lambda|Ric|^2)^2 - C. \tag{3.6}$$

Next, let $\varphi(t)$ be a smooth function on $\mathbb{R}^+$ so that $0 \leq \varphi(t) \leq 1$, $\varphi(t) = 1$ for $0 \leq t \leq R_0$, $\varphi(t) = 0$ for $t \geq 2R_0$, and

$$t^2\left(|\varphi'(t)|^2 + |\varphi''(t)|\right) \leq c \tag{3.7}$$

for some universal constant c and $R_0 > 0$ arbitrary large. We now take $\varphi = \varphi(F(x))$ as a cut-off function with support in $D(2R_0)$. Note that

$$|\nabla\varphi| = |\varphi'||\nabla F| \leq \frac{c}{R_0} \quad \text{and} \quad |\Delta_f\varphi| \leq |\varphi'\Delta_f F| + |\varphi''||\nabla F|^2 \leq \frac{c}{R_0} \tag{3.8}$$

on $D(2R_0) \setminus D(R_0)$ for some universal constant c.

Setting $u = |Rm| + \lambda|Ric|^2$ and $G = \varphi^2 u$, then direct computations, (3.6) and (3.8) yield

$$\begin{aligned}
\varphi^2\Delta_f G &= \varphi^4\Delta_f u + \varphi^2 u\Delta_f(\varphi^2) + 2\varphi^2\nabla u \cdot \nabla\varphi^2 \\
&\geq \varphi^4\left(u^2 - C\right) + \varphi^2 u\left(2\varphi\Delta_f\varphi + 2|\nabla\varphi|^2\right) + 2\nabla G \cdot \nabla\varphi^2 - 8|\nabla\varphi|^2 G \\
&\geq G^2 + 2\nabla G \cdot \nabla\varphi^2 - CG - C.
\end{aligned}$$

Now it follows from the maximum principle that $G \leq C$ on $D(2R_0)$ by some constant $C > 0$ depending on c_1 but independent of R_0. Hence $u = |Rm| + \lambda |Ric|^2 \leq C$ on $D(R_0)$. Since $R_0 > 0$ is arbitrary large, we see that

$$\sup_{x \in M} |Rm| \leq \sup_{x \in M} \left(|Rm| + \lambda |Ric|^2 \right) \leq C.$$

This completes the proof of Theorem 3.1. □

Proposition 3.2. *Let (M^4, g_{ij}, f) be a complete noncompact gradient steady soliton with positive Ricci curvature $Ric > 0$ and R attains its maximum at $x_0 \in M^4$. Then, function $u = \frac{|Rm| + \lambda |Ric|^2}{R}$, with $\lambda > 0$ sufficiently large, satisfies the differential inequality*

$$\Delta_f u \geq u^2 R - CR - 2\nabla u \cdot \nabla (\log R)$$

for some constant $C > 0$.

Proof. First of all, similar to deriving (3.4)-(3.6) in the proof of Theorem 3.1, by choosing λ sufficiently large we have

$$\begin{aligned} \Delta_f (|Rm| + \lambda |Ric|^2) &\geq (|Rm| + \lambda |Ric|^2)^2 - 4\lambda^2 |Ric|^4 - \lambda (|Ric|^4 + |Ric|^2) \\ &\geq (|Rm| + \lambda |Ric|^2)^2 - C|Ric|^2 \end{aligned}$$

for some constant $C > 0$. Here we have also used the fact (3.3).

Thus, by a direct computation,

$$\begin{aligned} \Delta_f u &= R^{-1} \Delta_f (|Rm| + \lambda |Ric|^2) + (uR) \Delta_f (R^{-1}) + 2\nabla (uR) \cdot \nabla (R^{-1}) \\ &\geq \frac{(|Rm| + \lambda |Ric|^2)^2 - C|Ric|^2}{R} + (uR) \left[2 \frac{|Ric|^2}{R^2} + 2 \frac{|\nabla R|^2}{R^3} \right] \\ &\quad - \frac{2}{R^2} \left(u |\nabla R|^2 + R \nabla u \cdot \nabla R \right) \\ &\geq Ru^2 - CR - 2\nabla u \cdot \nabla \log R. \end{aligned}$$

□

Theorem 3.2. *Let (M^4, g_{ij}, f) be a complete noncompact gradient steady Ricci soliton with $Ric > 0$ such that R attains its maximum. Suppose R has at most linear decay, $R(x) \geq c/r(x)$, outside a compact set. Then*

$$\sup_{x \in M} \frac{|Rm|}{R} \leq C.$$

Proof. Fix λ sufficient large so that Proposition 3.2 holds and set $u = \frac{|Rm| + \lambda |Ric|^2}{R}$. Next, let $\varphi(t)$ be a smooth function on $\mathbb{R}^+$ so that $\varphi(t) = \frac{d-t}{d}$ for $0 \leq t \leq d$, $\varphi(t) = 0$ for $t \geq d$. Let $\varphi = \varphi(F)$ and $G = \varphi^2 u$. Then,

$$\begin{aligned} |\nabla \varphi| &= |\varphi' \nabla F| \leq \frac{1}{d}, \\ \Delta_f \varphi &= \varphi' \Delta_f F = -\frac{1}{d}. \end{aligned}$$

Then, outside $D(1)$, we have

$$\begin{aligned}\varphi^2\triangle_f(G) &= \varphi^4(\triangle_f u)+\varphi^2 u(\triangle_f\varphi^2)+2\varphi^2\nabla\varphi^2\cdot\nabla u\\ &\geq \varphi^4\left(Ru^2-cR-2\nabla u\cdot\nabla\log R\right)\\ &\quad +2\left(\varphi\triangle_f\varphi+|\nabla\varphi|^2\right)G+2\varphi^2\nabla\varphi^2\cdot\nabla u\\ &\geq RG^2-cR+4\varphi\left(\nabla\varphi\cdot\nabla\log R\right)G\\ &\quad -\frac{8}{d}G+\left(2\nabla\varphi^2-2\varphi^2\nabla\log R\right)\cdot\nabla G.\end{aligned}$$

Now by (2.2), (3.2) and $Ric>0$, we have $|\nabla\log R|=2|\frac{Ric(\nabla f)}{R}|\leq 2$. Also, when R has at most linear decay outside some $D(t_0)$ and for $d>t_0$, we have $R\geq\frac{a}{d}$ in $D(d)\backslash D(t_0)$ for some constant $a>0$ independent d. Therefore there exists c independent of d so that,

$$\begin{aligned}\varphi^2\triangle_f(G) &\geq RG^2-cR-\frac{c}{d}G+\left(2\nabla\varphi^2-2\varphi^2\nabla\log R\right)\cdot\nabla G\\ &\geq \frac{1}{2}RG^2-cR+\left(2\nabla\varphi^2-2\varphi^2\nabla\log R\right)\cdot\nabla G.\end{aligned}$$

Therefore, it follows from stardard maximum principle argument that $u\leq C$ on M^4, hence $|Rm|\leq CR$ on M^4.

□

4. The proof of Theorem 1.2

In this section, we prove our second main result, Theorem 1.2 in the introduction. Throughout the section we assume (M^4,g_{ij},f) is a complete noncompact, non Ricci-flat, 4-dimensional gradient steady Ricci soliton such that

$$\lim_{x\to\infty}R(x)=0. \tag{4.1}$$

Note that, by Lemma 2.2 and Remark 2.1, (M^4,g_{ij},f) necessary has strictly positive scalar curvature $R>0$.

First of all, we need the following useful Laplacian comparison type result for gradient Ricci solitons.

Lemma 4.1. *Let (M^n,g_{ij},f) be any gradient steady Ricci soliton satisfying (1.1) and let $r(x)=d(x_0,x)$ denote the distance function on M^n from a fixed base point x_0. Suppose that*

$$Ric\leq(n-1)K$$

on the geodesic ball $B(x_0,r_0)$ for some constants $r_0>0$ and $K>0$. Then, for any $x\in M^n\setminus B(x_0,r_0)$, we have

$$\Delta_f r(x)\leq(n-1)\left(\frac{2}{3}Kr_0+r_0^{-1}\right).$$

Remark 4.1. Lemma 4.1 is a special case of a more general result valid for solutions to the Ricci flow due to Perelman [26], see, e.g., Lemma 3.4.1 in [10]. Also see [18] and [29] for a different version.

Theorem 4.1. *Let (M^4, g_{ij}, f), which is not Ricci-flat, be a complete noncompact gradient steady Ricci soliton. If $\lim_{x\to\infty} R(x) = 0$, then, for each $0 < a < 1$, there exists a constant $C > 0$ such that*

$$\sup_{x\in M} |Ric|^2 \le CR^a \qquad \text{and} \qquad \sup_{x\in M} |Rm| \le C.$$

Proof. The proof is similar to that of Munteanu-Wang [25] except we need to use the distance function to cut-off rather than the potential function since the potential function may not be proper.

Since $\lim_{x\to\infty} R(x) = 0$, it follows from (2.4) that

$$|\nabla f| \ge c_1 > 0$$

for some $0 < c_1 < 1$ outside a compact set. By Lemma 2.4 and Lemma 3.1, we have

$$\begin{aligned} \Delta_f |Ric|^2 &\ge 2|\nabla Ric|^2 - C|Rm||Ric|^2 \\ &\ge 2|\nabla Ric|^2 - C\left(|\nabla Ric| + |Ric|^2 + |Ric|\right)|Ric|^2. \end{aligned}$$

Also, since $R > 0$ on M^4, by using the first identity in Lemma 2.3 we have

$$\Delta_f \left(\frac{1}{R^a}\right) = 2a\frac{|Ric|^2}{R^{a+1}} + a(a+1)\frac{|\nabla R|^2}{R^{a+2}}.$$

Hence,

$$\begin{aligned} \Delta_f \left(\frac{|Ric|^2}{R^a}\right) &= \frac{\Delta_f |Ric|^2}{R^a} + |Ric|^2 \Delta_f \left(\frac{1}{R^a}\right) + 2\nabla|Ric|^2 \cdot \nabla\left(\frac{1}{R^a}\right) \\ &\ge \frac{2|\nabla Ric|^2}{R^a} - C\frac{\left(|\nabla Ric| + |Ric|^2 + |Ric|\right)|Ric|^2}{R^a} \\ &\quad + |Ric|^2\left[2a\frac{|Ric|^2}{R^{a+1}} + a(a+1)\frac{|\nabla R|^2}{R^{a+2}}\right] - 4a\frac{|Ric|\,|\nabla|Ric||\,|\nabla R|}{R^{a+1}} \end{aligned}$$

Apply Cauchy's inequality to the last term

$$\begin{aligned} -4a\frac{|Ric|\,|\nabla|Ric||\,|\nabla R|}{R^{a+1}} &\ge -4a\frac{|Ric|\,|\nabla Ric|\,|\nabla R|}{R^{a+1}} \\ &\ge -a(a+1)\frac{|Ric|^2|\nabla R|^2}{R^{a+2}} - \frac{4a}{a+1}\frac{|\nabla Ric|^2}{R^a}. \end{aligned}$$

Thus, we have

$$\begin{aligned} \Delta_f\left(|Ric|^2 R^{-a}\right) &\ge \frac{2(1-a)}{1+a}\frac{|\nabla Ric|^2}{R^a} - C\frac{|\nabla Ric||Ric|^2}{R^a} \\ &\quad - C\frac{|Ric|^4 + |Ric|^3}{R^a} + 2a\frac{|Ric|^4}{R^{a+1}} \\ &\ge (2a - \frac{CR}{1-a})\frac{|Ric|^4}{R^{a+1}} - C\frac{|Ric|^3}{R^a}. \end{aligned}$$

Therefore, for $u = \frac{|Ric|^2}{R^a}$, we have derived the differential inequality

$$\Delta_f u \ge (2a - \frac{CR}{1-a})u^2 R^{a-1} - Cu^{3/2}R^{a/2}. \tag{4.2}$$

By assumption (4.1), for any $0 < a < 1$, we can choose a fixed $d_0 > 0$ depending on a and sufficiently large so that

$$(2a - \frac{CR}{1-a}) \geq a \tag{4.3}$$

outside the geodesic ball $B(x_0, d_0)$.

Next, for any $D_0 > 2d_0$, we choose a function $\varphi(t)$ as follows: $0 \leq \varphi(t) \leq 1$ is a smooth function on $\mathbb{R}$ such that

$$\varphi(t) = \begin{cases} 1, & 2d_0 \leq t \leq D_0, \\ \\ 0, & t \leq d_0 \text{ or } t \geq 2D_0. \end{cases}$$

Also,

$$t^2|\varphi''(t)| \leq c \qquad \text{and} \qquad 0 \geq \varphi'(t) \geq -\frac{c}{D_0}, \text{ if } 2d_0 \leq t \leq 2D_0. \tag{4.4}$$

Now we use $\varphi = \varphi(r(x))$ as a cut-off function whose support is in $B(x_0, 2D_0) \setminus B(x_0, d_0)$. Note that, by (4.4) and Lemma 4.1, we get

$$|\nabla\varphi|^2 = |\varphi'|^2 \leq \frac{c}{D_0^2} \quad \text{and} \quad \Delta_f\varphi = \varphi'\Delta_f r(x) + \varphi'' \geq -\frac{C}{D_0}. \tag{4.5}$$

on $B(x_0, 2D_0) \setminus B(x_0, 2d_0)$ respectively.

Setting $G = \varphi^2 u$, then by our choice of φ and (4.5), we see that

$$\begin{aligned} \varphi^2\Delta_f G &= \varphi^4\Delta_f u + \varphi^2 u\Delta_f\varphi^2 + 2\varphi^2(\nabla u \cdot \nabla\varphi^2) \\ &\geq \varphi^4\left(au^2R^{a-1} - Cu^{3/2}R^{a/2}\right) + 2\varphi^2 u(\Delta_f\varphi^2) - 8|\nabla\varphi|^2 G + 2\nabla G \cdot \nabla\varphi^2 \\ &\geq aG^2R^{a-1} - CG^{3/2}R^{a/2} - CG + 2\nabla G \cdot \nabla\varphi^2. \end{aligned}$$

Assume G achieves its maximum at some point $p \in B(x_0, 2D_0)$. If $p \in B(x_0, 2D_0)\setminus B(x_0, 2d_0)$, then it follows from the maximum principle that

$$0 \geq aG^2(p)R^{a-1}(p) - CG^{3/2}(p)R^{a/2}(p) - CG(p).$$

On the other hand, noticing that the fact $0 < a < 1$ and R uniformly bounded from above, implies.

$$G(p) \leq C$$

for some constant C depending on a but independent of D_0.

Thus,

$$\max_{B(x_0, D_0)} u \leq \max_{B(x_0, 2D_0)} G \leq \max\left\{C, \max_{B(2d_0)} u\right\} \leq C'$$

for some $C' > 0$ indepedent of D_0. Therefore $|Ric|^2 \leq CR^a$ on M^4.

It remains to show $|Rm| \leq C$ on M^4. However, once we know $\sup_{x\in M} Ric \leq C$, $|Rm| \leq C$ follows essentially from the same argument as in the proof of Theorem 3.1. We leave the detail for the interested reader.

□

Lemma 4.2. *Let (M^4, g_{ij}, f), which is not Ricci-flat, be a complete noncompact gradient steady Ricci soliton with $\lim_{x\to\infty} R(x) = 0$. Then for each $0 < a < 1$ and $\mu > 0$, there exist constants $\lambda > 0$ and $D > 0$ so that function*

$$v = \frac{|Rm|^2 + \lambda|Ric|^2}{R^a}$$

satisfies the differential inequality

$$\Delta_f v \geq \mu v - D.$$

Proof. By Lemma 2.4 and Theorem 4.1,

$$\begin{aligned}\Delta_f v &= \frac{\Delta_f(|Rm|^2+\lambda|Ric|^2)}{R^a} + vR^a\Delta_f(\frac{1}{R^a}) + 2\nabla(vR^a)\cdot\nabla(R^{-a}) \\ &\geq \frac{2|\nabla Rm|^2+2\lambda|\nabla Ric|^2}{R^a} - c\frac{|Rm|^2+\lambda|Ric|^2}{R^a} \\ &\quad +(|Rm|^2+\lambda|Ric|^2)\left[-a\frac{\triangle_f R}{R^{a+1}} + a(a+1)\frac{|\nabla R|^2}{R^{a+2}}\right] \\ &\quad -4a\frac{|Rm||\nabla Rm||\nabla R|}{R^{a+1}} - 4a\lambda\frac{|Ric||\nabla Ric||\nabla R|}{R^{a+1}}.\end{aligned}$$

By using Cauchy's inequality to terms with $|\nabla R|$,

$$\begin{aligned}\Delta_f v &\geq \frac{2|\nabla Rm|^2+2\lambda|\nabla Ric|^2}{R^a} - c\frac{|Rm|^2+\lambda|Ric|^2}{R^a} \\ &\quad -\frac{4a}{a+1}\frac{|\nabla Rm|^2}{R^a} - \frac{4a\lambda}{a+1}\frac{|\nabla Ric|^2}{R^a} \\ &\geq \frac{2\lambda(1-a)}{1+a}\frac{|\nabla Ric|^2}{R^a} - c\frac{|Rm|^2+\lambda|Ric|^2}{R^a}.\end{aligned}$$

Now by Proposition 3.1, for some constant $\epsilon > 0$, we have

$$\begin{aligned}2\epsilon|Rm|^2 &\leq \left(|\nabla Ric| + |Ric|^2 + |Rc|\right)^2 \\ &\leq 2|\nabla Ric|^2 + 2(|Ric|^2+|Ric|)^2.\end{aligned}$$

Thus,

$$\begin{aligned}\Delta_f v &\geq \left[\frac{2\epsilon\lambda(1-a)}{1+a} - c\right]\frac{|Rm|^2}{R^a} - \left[2\lambda\frac{1-a}{1+a}(|Ric|+1)^2 + c\lambda\right]\frac{|Ric|^2}{R^a} \\ &\geq [\epsilon\lambda(1-a)-c]\,(v - \lambda\frac{|Ric|^2}{R^a}) - \lambda\left[2(1-a)(|Ric|+1)^2 + c\right]\frac{|Ric|^2}{R^a}.\end{aligned}$$

Therefore, by Theorem 4.1, for each $0 < a < 1$ and $\mu > 0$ one can choose $\lambda \geq C/(1-a)$, with $C > 0$ depending on μ and sufficiently large, so that

$$\Delta_f v \geq \mu v - D$$

for some constant $D > 0$ depending on λ. □

Theorem 4.2. *Let (M^4, g_{ij}, f), which is not Ricci-flat, be a complete noncompact gradient steady Ricci soliton with $\lim_{r\to\infty} R = 0$. Suppose R has at most polynomial decay, i.e., $R(x) \geq C/r^k(x)$ outside $B(r_0)$ for some fixed $r_0 > 1$, some constant $c > 0$ and positive integer k. Then, for each $0 < a < 1$, there exists a constant C such that*

$$|Rm| \leq CR^{a/2}.$$

Proof. Let $p=\frac{k}{2}$. Consider the following function on $\mathbb{R}^+$:

$$\varphi(t)=\begin{cases}\left(\frac{d-t}{d}\right)^p & 0\le t\le d\\ \\ 0 & t\ge d.\end{cases}$$

Next, let $\varphi=\varphi(r(x))$ on M^4. Then we have

$$\begin{aligned}|\nabla\varphi| &= \frac{p}{d}\left(\frac{d-r}{d}\right)^{p-1}|\nabla r|=\frac{p}{d-r}\varphi,\\ \triangle_f\varphi &= -\frac{p}{d}\left(\frac{d-r}{d}\right)^{p-1}\triangle_f r+\frac{p(p-1)}{d^2}\left(\frac{d-r}{d}\right)^{p-2}|\nabla r|^2\\ &= \left[-\frac{p}{d-r}\triangle_f r+\frac{p(p-1)}{(d-r)^2}\right]\varphi\end{aligned}$$

Consider $w=v-\frac{D}{\mu}$ with $v=\frac{|Rm|^2+\lambda|Ric|^2}{R^a}$, μ and D as in Lemma 4.2. Then, w satisfies

$$\triangle_f w\ge\mu w.$$

Let $G{=}\varphi^2 w$, then outside $B(r_0)$, we have

$$\begin{aligned}\triangle_f G &= (\Delta_f\varphi^2)w+\varphi^2\Delta_f w+2(\nabla\varphi^2)\cdot\nabla w\\ &\ge \left(2\varphi\Delta_f\varphi+2|\nabla\varphi|^2\right)w+\mu\varphi^2 w+4\varphi\nabla\varphi\cdot\nabla\frac{G}{\varphi^2}\\ &\ge \left(\mu+\frac{2\triangle_f\varphi}{\varphi}-6\frac{|\nabla\varphi|^2}{\varphi^2}\right)G+\frac{4}{\varphi}\langle\nabla G,\nabla\varphi\rangle. \qquad (4.6)\end{aligned}$$

Recall that $G=0$ outside $B(d)$. Now consider a maximum point q of G.

Case 1. $G(q)\le 0$. Then, $\max\limits_{B(d)} w\le 0$.

Case 2. $G(q)>0$ and $q\in B(r_0)$. Then, on $\Omega=B((1-\frac{1}{2^{1/p}})d)$, we have

$$\begin{aligned}\max_\Omega w &\le \max_\Omega\frac{1}{\varphi^2}\cdot G(q)\\ &\le 4G(q)\\ &\le 4\max_{B(r_0)} w.\end{aligned}$$

Case 3. $G(q)>0$ and $q\notin B(r_0)$. Then, by (4.6) and Lemma 4.1, at q we have

$$\begin{aligned}0 &\ge \mu+2\frac{\triangle_f\varphi}{\varphi}-6\frac{|\nabla\varphi|^2}{\varphi^2}\\ &\ge \mu-2pK_0\frac{1}{d-r}-(4p^2+2p)\frac{1}{(d-r)^2}\end{aligned}$$

for some constant $K_0>0$ depending on r_0 and $\max_{B(r_0)}|Ric|$. Hence $\frac{1}{d-r(q)}>C$ for some constant C depending on μ, $p=k/2$ and K_0. Thus, we have

$$d-r(q)\le c \qquad (4.7)$$

for some constant $c>0$ independent of d.

Therefore,

$$\begin{aligned}\max_{\Omega} w &\leq \max_{\Omega} \frac{1}{\varphi^2} \cdot G(q) \\ &\leq 4G(q) \\ &\leq 4\frac{(d-r(q))^{2p}}{d^{2p}} \frac{\left(|Rm|^2+\lambda|Ric|^2\right)}{R^a}(q) \\ &\leq C\frac{r^{ak}(q)}{d^{2p}} \\ &\leq Cd^{(a-1)k} \leq C\end{aligned}$$

for some constant $C > 0$ independent of d. Since $d > r_0$ is arbitrary, we obtain $\sup_M w \leq C$, and hence $|Rm|^2 \leq CR^a$ on M^4 for each $0 < a < 1$.

□

References

[1] Brendle, S., *Rotational symmetry of self-similar solutions to the Ricci flow*, Invent. Math. **194** (2013), no. 3, 731-764.

[2] Brendle, S., *Rotational symmetry of Ricci solitons in higher dimensions*, J. Differential Geom. **97** (2014), 191–214.

[3] Bryant, R., unpublished work (http://math.duke.edu/∼bryant/3DRotSymRicciSolitons.pdf)

[4] Buzano, M., Dancer, A. S., Gallaugher, M. and Wang, M., *A family of steady Ricci solitons and Ricci-flat metrics*, preprint (2014), arXiv:1309.6140v2 [math.DG].

[5] Cao, H.-D., *Existence of gradient Kähler-Ricci solitons, Elliptic and Parabolic Methods in Geometry* (Minneapolis, MN, 1994), A K Peters, Wellesley, MA, (1996) 1–16.

[6] Cao, H.-D., *Recent progress on Ricci solitons*, Recent advances in geometric analysis, 1–38, Adv. Lect. Math. (ALM), **11** *Int. Press, Somerville, MA*, 2010. MR2648937

[7] Cao, H.-D., Catino, G., Chen, Q., Mantegazza, C., and Mazzieri, L., *Bach-flat gradient steady Ricci solitons*, Calc. Var. Partial Differential Equations, **49** (2014), no. 1-2, 125-138.

[8] Cao H.-D. and Chen, Q., *On locally conformally flat gradient steady Ricci solitons*, Trans. Amer. Math. Soc., **364** (2012), 2377-2391.

[9] Cao H.-D. and Zhou, D., *On complete gradient shrinking Ricci solitons*, J. Differential Geom., **85** (2010), no. 2, 175–186.

[10] Cao H.-D. and Zhu, X.-P., *A complete proof of the Poincaré and geometrization conjectures—application of the Hamilton-Perelman theory of the Ricci flow*, Asian J. Math. **10** (2006), no. 2, 165–492. MR2233789 (2008d:53090)

[11] Catino, G. and Mantegazza, C., *Evolution of the Weyl tensor under the Ricci flow*, Ann. Inst. Fourier. **61** (2011), no. 4, 1407–1435.

[12] Chen, B.-L., *Strong uniqueness of the Ricci flow*, J. Differential Geom. **82** (2009), 363-382. MR2520796 (2010h:53095)

[13] Chen, X.X. and Wang, Y., *On four-dimensional anti-self-dual gradient Ricci solitons*, to appear in J. Geom. Anal.

[14] Chow, B. et al., *The Ricci flow: techniques and applications. Part I Geometric aspects.* Mathematical Surveys and Monographs, **135**. *American Mathematical Society, Providence, RI*, 2007. MR2302600 (2008f:53088)

[15] Chow, B., Lu, P. and Yang, B., *Lower bounds for the scalar curvatures of noncompact gradient Ricci solitons*, C. R. Math. Acad. Sci. Paris **349** (2011), no. 23-24, 1265–1267.

[16] Dancer, A. and Wang, M., *Some new examples of non-Kähler Ricci solitons*, Math. Res. Lett. **16** (2009), 349–363.

[17] Dancer, A. and Wang, M., *On Ricci solitons of cohomogeneity one*, Ann. Glob. Anal. Geom., **39** (2011) 259–292.

[18] Fang, F., Li, X.-D. & Zhang, Z., *Two Generalizations of Cheeger-Gromoll Splitting theorem via Bakry-Emery Ricci Curvature*, Ann. Inst. Fourier (Grenoble) **59** (2009), no. 2, 563–573.

[19] Feldman, M., Ilmanen, T. and and Knopf, D., *Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons*, J. Diff. Geom., **65** (2003), 169 –209.

[20] Fernndez-López, M. and Garca-Río, E., *A sharp lower bound for the scalar curvature of certain steady gradient Ricci solitons*, Proc. Amer. Math. Soc. **141** (2013), no. 6, 2145–2148.
[21] Hamilton, R. S., *The Ricci flow on surfaces*, Contemporary Mathematics **71** (1988), 237-261. MR0954419 (89i:53029)
[22] Hamilton, R. S., *The formation of singularities in the Ricci flow*, Surveys in Differential Geometry (Cambridge, MA, 1993), **2**, 7-136, International Press, Cambridge, MA, 1995. MR1375255 (97e:53075)
[23] Ivey, T., *New examples of complete Ricci solitons*, Proc. AMS **122** (1994), 241-245.
[24] Munteanu, O. and Sesum, N., *On gradient Ricci solitons*, J. Geom. Anal. **23** (2013), no. 2, 539–561.
[25] Munteanu, O. and Wang, J., *Geometry of shrinking Ricci solitons*, preprint (2014), arXiv:1410.3813v1.
[26] Perelman, G., *The entropy formula for the Ricci flow and its geometric applications*, arXiv:math.DG/0211159.
[27] Perelman, G., *Ricci flow with surgery on three manifolds*, arXiv:math.DG/0303109.v1
[28] Petersen, P. and Wylie, W., *Rigidity of gradient Ricci solitons*, Pacific J. Math, **241** (2009), 329–345.
[29] Wei, G. and Wylie, W., *Comparison Geometry for the Bakry-Emery Ricci Tensor*, J. Differential Geometry **83** (2009), 377–405.
[30] Wei, G. and Wu, P., *On volume growth of gradient steady Ricci solitons*, Pacific J. Math. **265** (2013), no. 1, 233–241.
[31] Wu, P., *On the potential function of gradient steady Ricci solitons*, J. Geom. Anal., **23** (2013), 221-228.

Department of Mathematics, University of Macau, Macao & Lehigh University, Bethlehem, PA 18015
E-mail address: huc2@lehigh.edu

Department of Mathematics, Lehigh University, Bethlehem, PA 18015
E-mail address: xic210@lehigh.edu